

Time to scale-up point-of-care ultrasound (POCUS) in obstetrics and gynaecology in Nigeria

Onyekpa IJ^{1,2,*}, Odugu BU¹, Okafor II^{1,2}, Ugwu IA¹, Onah LN^{1,2}, Nweze SO¹ and Nevo CO¹

¹ Department of Obstetrics and Gynaecology, Enugu State University of Science and Technology (ESUT) College of Medicine/ESUT Teaching Hospital, Parklane, Enugu, Nigeria.

² Part-time healthcare providers to Semino Hospital and Maternity, Abakpa Nike, Enugu, Nigeria.

World Journal of Advanced Pharmaceutical and Medical Research, 2024, 06(02), 001–006

Publication history: Received on 13 February 2024; revised on 17 April 2024; accepted on 20 April 2024

Article DOI: <https://doi.org/10.53346/wjapmr.2024.6.2.0033>

Abstract

The availability, affordability and portability of high-resolution-ultrasound machine in recent years have made ultrasound an indispensable point-of-care diagnostic and procedural tool in current practice of Obstetrics and Gynaecology. POCUS is a goal-directed (focused/emergency) ultrasound examination performed by the healthcare provider in the consulting office or patient's bedside to make a specific diagnosis that will direct timely and definitive treatment of the patient. It improves clinical outcomes, prevents hospital delays, promotes swift definitive treatment, reduces cost and enhances patient satisfaction without exposure to ionizing radiation. It is a readily available care in Obstetrics and Gynaecology. Skills acquisition in POCUS should be incorporated in teaching programs for midwives, medical students, resident doctors and consultants. This review aims to summarize the literature on the advances in ultrasound technology, indications and practice guidelines for clinicians' POCUS examinations in Obstetrics and Gynaecology. Recommendations to fill the gaps in the current training and skills acquisition in Nigeria are proposed.

Keywords: POCUS; Obstetrics and Gynaecology; Indications; Guidelines; Scale-up-practice

1. Introduction

1.1. Point Of Care Ultrasound (POCUS)

POCUS can be defined as a goal-directed bedside or office clinician-performed ultrasound examination (Figure 1) for a specific diagnostic and/or interventional purpose [1, 2]. The main objective of POCUS during patient evaluation is to make a prompt diagnosis that will direct swift definitive treatment of the patient. It is a rapid but limited study performed by the same clinician who knows the patient's history and has performed the necessary physical examination. It reduces delays in diagnosis and initiation of necessary treatment [3]. The immediate diagnoses and swift definitive interventions are particularly relevant in obstetric and gynaecological emergencies where obstetric delays are known to aggravate adverse fetomaternal outcomes [4]. It is not, however, a substitute for an in-depth diagnostic ultrasound scan by skilled radiologists, but a complementary diagnostic tool that can make the clinician more efficient in medical/clinical decision-making. It can also be called emergency, focused or clinician-performed ultrasound.

* Corresponding author: Onyekpa IJ



Figure 1 Antenatal clinic office POCUS at 37 weeks gestational age in Semino Hospital and Maternity, Enugu to exclude placenta praevia in a known low-lying placenta at 18 weeks

2. Advances in ultrasound technology and relevance in clinical practice

Karl Theodore Dussik (Figure 2), an Austrian psychiatrist and neurologist, was the first physician to use a primitive-gigantic-complex ultrasound device to interrogate and visualize cerebral ventricles and brain tumors in 1946 [5]. Advances in ultrasound technology evolved from the ‘immersion-tank ultrasound systems’ to A-mode in the late 1950s, B-mode in the 1970s [6], real-time and gray-scale images in the 1980s [7], Doppler and color Doppler in the 1990s [8, 9]; and three dimensional and four-dimensional ultrasounds in the 2000s [10]. The use of artificial intelligence in ultrasound, new visualization methods and improved ergonomics are where the advances in ultrasound technology are currently going to [11].

The development of portable ultrasound devices including miniature handheld ones (Figure 3) in recent years has made it an indispensable point-of-care diagnostic and procedural tool in current practice of Obstetrics and Gynaecology. Ian Donald pioneered the use of obstetrics and gynecologic ultrasound on diagnosis of ovarian cyst [6]. POCUS has revolutionized the practice of medicine from the primitive bedside tools like the reflex hammer and stethoscope to bedside ultrasound that can make clinicians actually visualize immediately what they have palpated or auscultated. It is an extension of the physical examination and the greatest advancement in bedside diagnosis since the advent of stethoscope about two centuries ago. It is relevant in virtually every field of medicine and surgery. The integration of POCUS into routine clinical practice was, however, too slow in spite of its enormous positive impacts in clinical outcomes. It reduces failure and complication rates during procedures [12], rapidly narrows differential diagnoses [13], shortens times to definitive treatment [14], lowers cost of care [15], and reduces the exposure to ionizing radiation [16]. It involves patients in their case managements, enhances patient-doctor relationships and patient satisfactions [17]. POCUS should be a ‘must-know’ basic skill before a resident doctor qualifies to participate in the Part 1 Fellowship Examination in Obstetrics and Gynaecology in Nigeria.




Figure 2 karl Theodore dussik and his first medical ultrasound device in 1946 [5]



Figure 3 Handheld "miniature" ultrasound system for POCUS [11]

3. Indications of POCUS in Obstetrics and Gynaecology [18-23]

The thermal index (TI) and mechanical index (MI) effects of ultrasound were observed in the 1920s to kill fish that were within the ultrasonic beam [18]. These bio-effects of ultrasound were initially used for therapeutic applications. Later ultrasound was used to visualize internal anatomy [19]. There is a wide range of indications for POCUS (Table1) that one cannot imagine practicing modern Obstetrics and Gynaecology without basic ultrasound skills [20-22]. The impacts on clinical decision-makings are enormous [23]. These impacts are most beneficial when there are appropriate indications and the clinicians are competent to avoid misuse of ultrasound examinations that can lead to irrelevant findings, unnecessary further investigations, patient anxiety and even harmful treatments [23].

3.1. Indications for POCU

Gynaecological and first trimester ultrasound examinations [23, 24]:

- Vaginal bleeding
- Abdominal and pelvic pain
- Abdominal and pelvic masses
- Amenorrhoea
- Confirmation of an intrauterine pregnancy (IUP)
- Dating of pregnancy with crown–rump length of the embryo
- Monitor embryo and fetal growth and/or demise
- Confirmation of complete or incomplete abortion
- Diagnosis of ectopic pregnancy especially with TVS
- Location of missing contraceptive devices
- Measure endometrial thickness
- Detection of uterine polyps and fluid in the cavity
- Diagnosis of ovarian cysts and torsion
- Diagnosis of multiple pregnancy and the determinations of chorionicity and amnioicity.
- Serial measurements of cervical length for ultrasound indicated/directed cervical cerclage to avoidance unnecessary routine history indicated cerclage.
- Others include:
- Suspicion of uterine malformation
- Follicular tracking and baby sex selection

Procedural or interventional indications

- Ovum retrieval and embryo transfer
- Retrieval of missing intrauterine contraceptive device.
- Medical treatment of ectopic pregnancy with potassium chloride and methotrexate
- Aspiration of ovarian cysts

Second and third trimesters, labour and puerperium POCUS indications

- Assessment of fetal well-being: Fetal growth, macrosomia and intrauterine growth restriction
- Assessment of fetal lie, presentation, position and weight
- Placental localization
- Amniotic fluid volume estimation
- Accurate determination of presentation, position, and cervical dilation during labour and delivery management [24].
- Rupture of the pregnant uterus and free peritoneal fluid
- Puerperal uses to diagnosis haemoperitoneum, pelvic abscess, retained fragment of the placenta and missing abdominal mops or surgical instrument.

4. Training Curriculum, Practice Guidelines, and Skill Acquisition programs [25-30].

The availability of competent healthcare providers with basic ultrasound skills is critical for the integration of POCUS services into routine clinical practice [11, 21, 25-28]. The World Health Organization (WHO) [29], International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) [26] and American Institute of Ultrasound in Medicine (AIUM) [27] expert opinions have recognized the impact and importance of decentralization of POCUS service delivery through appropriate healthcare provider's training and task sharing. ISUOG and AIUM have developed consensus-based curriculum, standardized practice guidelines and competency assessment tools as modified in tables 2 and 3 that can be used as the minimum requirement for competency by training institutions [26, 27]. Obtaining 3 ultrasound images with good image optimizations and at the appropriate planes within 1 minutes for a specified task can be used as the minimum standards for attaining competency. The use of ultrasound mannequins during training will enhance the competency-assessment process.

4.1. Basic training curriculum [28] and practical batches of 3 students for 3 hours each day for 5 days

- Basic principles of ultrasound physics
- Knobology: frequency, resolution, depth gain, time gain compensation etc
- Ultrasound modes (B-mode, M-mode, Doppler, two-dimensional (2D) and three-dimensional (3D)).
- Ultrasound transducers: types and applications.
- Image orientations: Trans Abdominal Ultrasound (TAS) and Trans Vaginal Ultrasound (TVS)
- Bio-effects of ultrasound (As Low As Reasonably Achievable (ALARA) principle) [30].
- Patient review: identification, counseling, indications, position and inputting information.
- Ultrasound examination (Hand-on-training in groups of not more than 3): coupling agents, ergonomic practices, correct transducer manipulation and image orientation, image labeling and storage and documentation of findings. Careful transducer handling, cleaning and disinfection.
- Recommended videos to watch during breaks
 - The Principles and Physics of Ultrasound Imaging: Medical Aid Films - Films For Life
 - Made in partnership with ISUOG,
 - Ultrasound Machine Basics- Knobology, Probes, and Modes www.pocus101
 - Gynecology/Pelvic Ultrasound Made Easy: Step-By-Step Guide www.pocus101.com
 - The Basic Steps of an Obstetric Ultrasound Examination: Medical Aid Films - Films For Life. Made in partnership with ISUOG,
 - ISUOG: Plan your movement and improve your image: Dr. Hisham Mirghani

4.2. Basic first trimester ultrasound examination

Demonstration of steps for performance of first-trimester trans-abdominal and trans-vaginal ultrasound examinations. Obtaining 3 images of each of the following with good image optimizations within 1 minute can be accepted as minimum competency criteria during the training.

- Ultrasound examination of non-pregnant uterus during various menstrual cycle phases, normal ovary, simple cyst and corpus lutein
- Demonstrations of intrauterine gestational sac, yolk sac, amnion and number of embryo/fetuses.
- Documentation of embryo/fetus cardiac activity or death. Diagnosis of missed, incomplete abortions and molar pregnancy.
- Pregnancy dating in first trimester especially with CRL.
- Ultrasound evaluation of ectopic pregnancy, free peritoneal fluid and pregnancy of unknown origin.
- .

4.3. Senior Training of 5 days for 3 hours each day.

Ultrasound in the second and third trimesters

- Components of basic second- and third-trimester ultrasound examinations.
- Fetal biometry: BPD, HC, AC, FL, macrosomia and fetal growth restriction.
- Placental localization, praevia, abruption, vasa praevia
- Ultrasound evaluation of twin gestations including chorionicity and amnionity.
- Amniotic fluid assessment: Oligohydramnios, Polyhydramnios and AFI
- Fetal anomaly scan
- TVS measurement of cervical length in second and third trimesters of pregnancy.
- Anatomic locations of leiomyomas, adenomyosis. and adnexal masses in pregnancy and labour
- Sonographic characteristics of polycystic ovaries and adnexal cyst torsion.
- Normal fetal anatomy and common malformations.
- Umbilical artery Doppler studies in evaluation of fetal growth restriction, TTTS and Doppler scan in evaluation of adnexal masses
- Writing the ultrasound report

5. Conclusion

Pocus is a 'must know-adjunct' skill to the routine history and physical examination in modern obstetric and gynaecological practice. The skill should be acquired through scaling-up of appropriate clinician's training curriculum, practice guidelines and task sharing. The positive impacts of POCUS on improving clinical outcomes are enormous and can no longer be ignored.

Compliance with ethical standards

Acknowledgments

We acknowledge the management and staff of SEMINO Hospital, Enugu for allowing us the use their POCUS facilities for this study.

Disclosure of conflict of interest

There was no conflict of interest in the course of the study

References

- [1] Fröhlich E, Beller K, Muller R, Herrmann M, Debove I, Klinger C et al (2020): Point of care ultrasound in geriatric patients: prospective evaluation of a portable handheld ultrasound device. *Ultraschall Med Eur J Ultrasound* 41(03):308–316. <https://doi.org/10.1055/a-0889-8070>
- [2] Sporea I (2014): Ultrasound for everyone – is it the right time? *Med Ultrason* 16(1):3–4
- [3] Moore CL, Copel JA (2011): Point-of-care ultrasonography. *N Engl J Med* 364(8):749–757
- [4] Vinita Das, Smriti Agrawal, Anjoo Agarwal (2009): Consequences of delay in obstetric care for maternal and perinatal outcomes. *International Journal of Gynaecology and Obstetrics* 109 (1):72-3. DOI: 10.1016/j.ijgo.2009.11.003
- [5] Frentzel-Beyme B. Vom Echolotzur, Farb doppler sonographie. *Der Radiologe*. April 2005;45(4):363–370.
- [6] Donald I, Macvicar J, Brown TG. Investigation of abdominal masses by pulsed ultrasound. *Lancet* 1958; 1:1188-1195.
- [7] Timor-Tritsch IE, Farine D, Rosen MG. A close look at early embryonic development with the high-frequency transvaginal transducer. *Am J Obstet Gynecol* 1988;159:676-681.
- [8] Nimura Y. Introduction of the ultrasonic Doppler technique in medicine: a historical perspective. *J Med Ultrasound* 1998;6:5-13.
- [9] FitzGerald DE, Drumm JE. Non-invasive measurement of human fetal circulation using ultrasound: a new method. *Br Med J* 1977;2:1450-1451

- [10] Timor-Tritsch IE, Monteagudo A. Three and four-dimensional ultrasound in obstetrics and gynecology. *Curr Opin Obstet Gynecol* 2007;19:157-175.
- [11] Jacques S Abramowicz (2021): Obstetric ultrasound: where are we and where are we going? <https://doi.org/10.14366/usg.20088>, pISSN: 2288-5919 • eISSN: 2288-5943, *Ultrasonography* 2021;40:57-74
- [12] Kanji HD, McCallum J, Sirounis D, MacRedmond R, Moss R, Boyd JH. Limited echocardiography-guided therapy in subacute shock is associated with change in management and improved outcomes. *J Crit Care*. 2014;29(5):700-705.
- [13] Al Deeb M, Barbic S, Featherstone R, Dankoff J, Barbic D. Point-of-care ultrasonography for the diagnosis of acute cardiogenic pulmonary edema in patients presenting with acute dyspnea: a systematic review and meta-analysis. *Acad Emerg Med*. 2014;21(8):843-852.
- [14] Melniker LA, Leibner E, McKenney MG, Lopez P, Briggs WM, Mancuso CA. Randomized controlled clinical trial of point-of-care, limited ultrasonography for trauma in the emergency department: the first sonography outcomes assessment program trial. *Ann Emerg Med*. 2006;48(3):227-235.
- [15] Parker L, et al. Musculoskeletal imaging: Medicare use, costs, and potential for cost substitution. *J Am Coll Radiol*. 2008;5(3):182-188.
- [16] Smith-Bindman R, et al. Ultrasonography versus computed tomography for suspected nephrolithiasis. *N Engl J Med*. 2014;371:1100-1110.
- [17] Howard ZD, et al. Bedside ultrasound maximizes patient satisfaction. *J Emerg Med*. 2014;46 (1):46-53.
- [18] Langevin P. Les ondes ultrasonores. *Rev Gen Electr* 1928;23: 626-634.
- [19] Gohe H, Wedekind T. Der Ultraschall in der Medizin. *Klin Wochenschr* 1940;19:25-29.
- [20] Abramowicz JS. Benefits and risks of ultrasound in pregnancy. *Semin Perinatol* 2013;37:295-300.
- [21] Recker, F., Weber, E., Strizek, B., Gembruch, U., Westerway, S. C., & Dietrich, C. F. (2021). Point-of-care ultrasound in obstetrics and gynecology. *Archives of gynecology and obstetrics*, 303(4), 871–876. <https://doi.org/10.1007/s00404-021-05972-5>
- [22] Reynolds TA, Amato S, Kulola I, Chen CJJ, Mfinanga J, Sawe HR (2018) Impact of point of-care ultrasound on clinical decision-making at an urban emergency department in Tanzania. *PLoS ONE* 13(4): e0194774. <https://doi.org/10.1371/journal.pone.0194774>
- [23] Lil Valentin (1999) Editorial: High-quality gynecological ultrasound can be highly beneficial, but poor-quality gynecological ultrasound can do harm. *Ultrasound Obstet Gynecol* 1999;13:1–7
- [24] Mohammed Algodí, Diana S. Wolfe and Cynthia C. Taub Commentry: The Utility of Maternal Point of Care Ultrasound on Labor and Delivery Wards *Journal of Cardiovascular Development and Disease*
- [25] Salomon LJ, Alfirevic Z, da Silva Costa F, Deter RL, Figueras F, Ghi T, Glanc P, Khalil A, Lee W, Napolitano R, Papageorghiou AT, Sotiriadis A, Stirnemann J, Toi A, Yeo G. ISUOG Practice Guidelines: ultrasound assessment of fetal biometry and growth. *Ultrasound Obstet Gynecol* 2019; 53: 715–723.
- [26] Bhide A, Acharya G, Baschat A, Bilardo CM, Brezinka C, Cafici D, Ebbing C, Hernandez-Andrade E, Kalache K, Kingdom J, Kiserud T, Kumar S, Lee W, Lees C, Leung KY, Malinger G, Mari G, Prefumo F, Sepulveda W, Trudinger B. ISUOG Practice Guidelines (updated): use of Doppler velocimetry in obstetrics. *Ultrasound Obstet Gynecol* 2021; 58: 331–339.
- [27] AIUM–ACR–ACOG–SMFM–SRU Practice Parameter for the Performance of Standard Diagnostic Obstetric Ultrasound Examinations. *Ultrasound Med*. 2018, 37, E13–E24. [CrossRef] [PubMed]. <https://www.aium.org/guidelines/obstetric>
- [28] A. Abuhamad, K. K. Minton, C. B. Benson, T. Chudleigh, L. Crites, P. M. Doubilet, R. Driggers, W. Lee, K. V. Mann, J. J. Perez, N. C. Rose, L. L. Simpson, A. Tabor And B. R. Benacerraf (2018): Obstetric and gynecologic ultrasound curriculum and competency assessment in residency training programs: consensus report.
- [29] Training in diagnostic ultrasound: essentials, principles and standards: report of a WHO study group (WHO technical report series; 875)
- [30] American Institute of Ultrasound in Medicine. AIUM official statement. As Low As Reasonably Achievable (ALARA) Principle. American Institute of Ultrasound in Medicine: Laurel, MD, 2014. <http://www.aium.org/officialStatements/39>